LAMPIRAN

Correlations

		Y	x1	x2	x3
Pearson Correlation	Y	1.000	.856	.148	.118
	x1	.856	1.000	-.075	.225
	x2	.148	-.075	1.000	.086
	x3	.118	.225	.086	1.000
Sig. (1-tailed)	Y	.	.000	.166	.220
	x1	.000	.	.312	.069
	x2	.166	.312	.	.288
	x3	.220	.069	.288	.
N	Y	45	45	45	45
	x1	45	45	45	45
	x2	45	45	45	45
	x3	45	45	45	45

Descriptive Statistics

	Mean	Std. Deviation	N
Y	1.32E4	13966.610	45
x1	7.3059E2	714.05763	45
x2	.4089	.18862	45
x3	1.7199082 E1	.90413494	45

Variables Entered/Removed[b]

Model	Variables Entered	Variables Removed	Method
1	x3, x2, x1[a]	.	Enter

a. All requested variables entered.

b. Dependent Variable: y

Model Summary[b]

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Durbin-Watson
1	.888[a]	.789	.773	6648.480	1.749

a. Predictors: (Constant), x3, x2, x1

b. Dependent Variable: y

ANOVA[b]

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	6.771E9	3	2.257E9	51.058	.000[a]
	Residual	1.812E9	41	4.420E7		
	Total	8.583E9	44			

a. Predictors: (Constant), x3, x2, x1

b. Dependent Variable: y

Coefficients[a]

Model		Unstandardized Coefficients		Standardized Coefficients	T	Sig.	Collinearity Statistics	
		B	Std. Error	Beta			Tolerance	VIF
1	(Constant)	13576.881	26707.932		.508	.614		
	x1	22.347	2.911	.790	7.676	.000	.924	1.082
	x2	23425.728	9626.327	.252	2.434	.019	.910	1.099
	x3	-1400.102	1614.106	-.091	-.867	.391	.897	1.115

a. Dependent Variable: y

One-Sample Kolmogorov-Smirnov Test

		Unstandardized Residual
N		45
Normal Parameters[a]	Mean	.0000000
	Std. Deviation	8.84852330E3
Most Extreme Differences	Absolute	.166
	Positive	.166
	Negative	-.153
Kolmogorov-Smirnov Z		1.117
Asymp. Sig. (2-tailed)		.165

a. Test distribution is Normal.







Apabila sampel data di SQRT maka diperoleh hasil sebagai berikut, bisa dibandingkan hasilnya dengan data yang diolah dengan data mentah tanpa di SQRT diatas.

Scatterplot



Normal P-P Plot of Regression Standardized Residual